Hydrobiologia

The International Journal of Aquatic Sciences

Volume 541, 1 June 2005

Primary Research Papers

Microhabitat type selection of caddisfly larvae (Insecta: Trichoptera) in a shallow lowland stream
G. Urbanič, M.J. Toman, C. Krušnik 1–12

Decomposition and mineralization of *Eichhornia crassipes* litter under aerobic conditions with and without bacteria
N.P.D. Gamage, T. Asaeda 13–27

Relationships between cyanobacterial production and the physical and chemical properties of a Midwestern Reservoir, USA
S.-H. Wang, A.R. Dzialowski, J.O. Meyer, F. deNoyelles, Jr., N.-C. Lim, W.W. Spotts, D.G. Huggins 29–43

Patterns of macroinvertebrate abundance in inland saline wetlands: a trophic analysis
E.A. Hart, J.R. Lovvorn 45–54

Ciliated protozoan communities and saprobic evaluation of water quality in the hilly zone of some tributaries of the Po River (northern Italy)
P. Madoni 55–69

Structural and functional phytoplankton responses to nutrient impoverishment in mesocosms placed in a shallow eutrophic reservoir (Garças Pond), São Paulo, Brazil
L.O. Crossetti, C.E.d.M. Bicudo 71–85

Distribution and toxicity of a new colonial *Microcystis aeruginosa* bloom in the San Francisco Bay Estuary, California
P.W. Lehman, G. Boyer, C. Hall, S. Waller, K. Gehrts 87–99

Coral mass- and split-spawning at a coastal and an offshore Venezuelan reefs, southern Caribbean
C. Bastidas, A. Cróquer, A.L. Zubillaga, R. Ramos, V. Kortnik, C. Weinberger, L.M. Márquez 101–106

Description of a new species from Laguna Chichancanab, Yucatan, Mexico: *Cyprinodon suavium* (Pisces: Cyprinodontidae)
U. Strecker 107–115

The importance of lake morphometry and catchment characteristics in limnology – ranking based on statistical analyses
L. Håkanson 117–137

Tadpole shrimp structure macroinvertebrate communities in playa lake microcosms
S.H. Yee, M.R. Willig, D.L. Moorhead 139–148

Bioavailability of zinc in the sediment to the estuarine amphipod *Grandidierella japonica*
J. Han, D. Ma, X. Quan, J. Wang, Q. Yan 149–154

Recovery from acidification in boreal lakes inferred from macroinvertebrates and subfossil chironomids
J. Hynynen, J.J. Meriläinen 155–173

Aquatic macroinvertebrate assemblages in mitigated and natural wetlands
C.K. Balcombe, J.T. Anderson, R.H. Fortney, W.S. Kordek 175–188

Biological and ecological features of saline lakes in northern Tibet, China
Z. Wen, Z. Mianping, X. Xianzhong, L. Xi-Fang, G. Ganlin, H. Zhihui 189–203

Veligers from the nudibranch *Dendronotus frondosus* show shell growth and extended planktonic period in laboratory culture
C.G. Sisson 205–213

Variation in stable isotope signatures of seston and a zooplanktivorous fish in a eutrophic Chinese lake
J. Xu, P. Xie, M. Zhang, H. Yang 215–220

The occurrence of a bloom-forming green alga *Pleodorina indica* (Volvocales) in the downstream reach of the River Malše (Czech Republic)
P. Znachor, J. Jezberová 221–228

Toward using *Margaritifera falcata* as an indicator of base level nitrogen and carbon isotope ratios: insights from two California Coast Range rivers
J.K. Howard, K.M. Cuffey, M. Solomon 229–236

Short Research Notes

Does the process of drying submerged macrophytes affect community structure and composition of epiphytic diatoms?
A. Vogel, T. Beier, J. Braun, U. Raeder 237–240

Benefit from an invader: American slipper limpet *Crepidula fornicata* reduces star fish predation on basibiont European mussels
D.W. Thieltges 241–244

Available online
www.springerlink.com